

## ***Drug Facts***

### ***Active ingredient***

Brimonidine tartrate (0.025%).....

### ***Purpose***

Redness reliever

### ***Use***

- relieves redness of the eye due to minor eye irritations

### ***Warnings***

**For external use only**

**Do not use** ■ if solution changes color or becomes cloudy

**Stop use and ask a doctor if** ■ you experience eye pain, changes in vision, continued redness or irritation of the eye

■ condition worsens or persists for more than 3 days

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### ***Directions***

- adults and children 5 years of age and over:
  - instill 1 drop in the affected eye(s) every 6-8 hours
  - do not use more than 4 times daily
  - remove contact lenses before use
  - wait at least 10 minutes before re-inserting contact lenses after use
- if using other ophthalmic products while using this

## ***Drug Facts*** (continued)

product, wait at least 5 minutes between each product

- to avoid contamination, do not touch tip of container to any surface
- replace cap after each use
- children under 5 years of age: consult a doctor

### ***Other information***

- store at 15°-25°C (59°-77°F)

### ***Inactive ingredients***

benzalkonium chloride, boric acid, calcium chloride dihydrate, glycerin, potassium chloride, sodium borate decahydrate, sodium chloride, water for injection. Hydrochloric acid and/or sodium hydroxide may be used to adjust pH.

### ***Questions or comments?***

☎ Call: 1-800-553-5340