

Drug Facts

Active ingredient

Brimonidine tartrate (0.025%).....

Purpose

Redness reliever

Use

- relieves redness of the eye due to minor eye irritations

Warnings

For external use only

Do not use ■ if solution changes color or becomes cloudy

Stop use and ask a doctor if ■ you experience eye pain, changes in vision, continued redness or irritation of the eye

■ condition worsens or persists for more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 5 years of age and over:
 - instill 1 drop in the affected eye(s) every 6-8 hours
 - do not use more than 4 times daily
 - remove contact lenses before use
 - wait at least 10 minutes before re-inserting contact lenses after use
- if using other ophthalmic products while using this

Drug Facts (continued)

product, wait at least 5 minutes between each product

- to avoid contamination, do not touch tip of container to any surface
- replace cap after each use
- children under 5 years of age: consult a doctor

Other information

- store at 15°-25°C (59°-77°F)

Inactive ingredients

benzalkonium chloride, boric acid, calcium chloride dihydrate, glycerin, potassium chloride, sodium borate decahydrate, sodium chloride, water for injection. Hydrochloric acid and/or sodium hydroxide may be used to adjust pH.

Questions or comments?

☎ Call: 1-800-553-5340